

Patient & Product Information

(Choline Salicylate 8.7% w/v, Lignocaine Hydrochloride 2% w/w, &
Benzalkonium Chloride 0.01% w/w)
Oral Pain Relief Gel

Active Ingredients:

Choline Salicylate8.7% w/w
Lignocaine Hydrochloride2.0% w/w
Benzalkonium Chloride0.01% w/w

Read this Patient Information before you start using this Oral Pain Relief Gel and each time you get a refill. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is this Gel?

This gel is a topical oral medication used to provide temporary relief from:

- Pain, swelling, and inflammation in the mouth caused by ulcers, sores, or other minor irritations.
- Discomfort associated with teething in infants and toddlers (as directed by a healthcare provider).
- Gum pain or irritation caused by dentures or orthodontic appliances.

Product Category:

- Oral Analgesic and Antiseptic Gel

Who should not use this Gel?

Do not use this if you:

- If you are allergic to choline salicylate, lignocaine hydrochloride, benzalkonium chloride, or any of the other ingredients in the gel.
- In children under 2 years unless prescribed by a healthcare provider.
- If you have a history of allergic reactions to salicylates or local anesthetics.

What should I tell my healthcare provider before using this product?

Before using this product, inform your healthcare provider if you:

- Have or have had bleeding disorders, allergies, or gum diseases.
- Are taking blood-thinning medications, aspirin, or other salicylates.
- Are pregnant, planning to become pregnant, or breastfeeding.
- Are currently using other topical or oral medications.

How should I use this Gel?

Use this gel exactly as directed by your healthcare provider or as indicated on the package.

- **For adults and children:** apply a small amount of gel (approximately a pea-sized quantity) to the affected area and gently massage.
- **For infants and toddlers:** apply a very small amount of gel to the gums using a clean finger or a cotton swab.
- Do not use this gel more than 3-4 times daily unless directed by your healthcare provider.
- Avoid eating or drinking immediately after application to allow the gel to take effect.

What are the possible side effects of This Gel?

The most common side effects include:

- Mild burning or tingling sensation at the application site.
- Temporary numbness or loss of sensation.

Rare but serious side effects include

- Allergic reactions such as rash, itching, swelling, or difficulty breathing.
- Prolonged or severe numbness.
- Tell your healthcare provider if you experience any side effects that bother you or do not go away.

How should I store this Gel?

- Store this gel at room temperature, between 20°C to 25°C (68°F to 77°F).
- Keep the tube tightly closed when not in use.
- Keep this gel and all medications out of the reach of children.

General information about the safe and effective use of this gel:

- Do not use this gel for a condition for which it was not prescribed.
- Do not share this gel with others, even if they have the same symptoms.
- You can ask your pharmacist or healthcare provider for more information.

What are the ingredients in this Gel?

Active ingredients:

- Choline Salicylate: Reduces pain and inflammation.
- Lignocaine Hydrochloride: Provides local anesthesia to numb the affected area.
- Benzalkonium Chloride: Acts as an antiseptic to prevent infections.

Inactive Ingredient's:

- Gel base