

Patient & Product Information

(Lignocaine 4% w/w, Metronidazole 1% w/w & Sucralfate 7% w/w Cream) **Topical Anesthetic, Antibacterial & Healing Agent**

Active Ingredients:

Lignocaine4.0% w/w
Metronidazole.....1.0% w/w
Sucralfate7.0% w/w

Read this Patient Information before you start using this Medicated Cream and each time you get a refill. This information does not replace advice from your healthcare provider about your medical condition or treatment.

What is this Medicated Cream?

This cream is a multi-functional topical medication used for:

- Relieving pain, burning, and discomfort in anorectal or dermatological conditions
- Treating localized bacterial infections, especially anaerobic infections
- Promoting wound healing in anal fissures, minor cuts, or skin abrasions
- Soothing inflamed mucosa and skin in perianal disorders

Who should not use this Medicated Cream?

Do not use this cream if you:

- Are allergic to Lignocaine, Metronidazole, Sucralfate, or any of the cream's ingredients
- Have severe liver dysfunction or blood disorders (without medical supervision)
- Are applying to deep, open wounds or infected skin unless prescribed
- Are pregnant or breastfeeding, unless advised by your healthcare provider

Who can use this Medicated Cream?

Adults and Elderly Patients:

- Safe when used as directed by a healthcare provider

Children (12 years and older):

- Use only under adult supervision; apply only as directed

Children under 12 years:

- Not recommended unless prescribed by a healthcare provider

What should I tell my healthcare provider before using this Medicated Cream?

Inform your doctor if you:

- Have a history of allergies or sensitive skin
- Have liver or kidney issues
- Are pregnant, planning pregnancy, or breastfeeding
- Are using other topical medications or antibiotics

How should I use this Medicated Cream?

- Use this cream exactly as prescribed by your healthcare provider.

For Adults and Children 12 Years and Older:

- Clean and dry the affected area before applying
- Apply a thin layer to the affected region 2–3 times daily, or as directed
- Wash hands after application (unless treating hands)
- Do not bandage unless directed
- Avoid contact with eyes, mouth, or mucous membranes

For Children Under 12 Years:

- Use only if prescribed and follow specific instructions

- Possible Side Effects of this Medicated Cream

Common Side Effects:

- Mild burning, stinging, or redness at the application site
- Temporary numbness or tingling in the treated area

Rare but Serious Side Effects:

- Allergic reactions (rash, swelling, itching, breathing difficulty)
- Neurological symptoms like dizziness or confusion (in case of accidental ingestion or large application areas)

In Children:

- Increased risk of systemic absorption, which may cause unwanted effects
- Stop using the cream and contact your doctor if side effects persist or worsen.

How should I store this Medicated Cream?

- Store at room temperature (20°C to 25°C / 68°F to 77°F)
- Do not freeze
- Keep away from heat and direct sunlight
- Keep out of reach of children

General Information for Safe Use:

- For external use only
- Do not use for any condition other than prescribed
- Do not share with others, even with similar symptoms
- Discontinue use if irritation persists or new symptoms appear
- For professional advice, consult your pharmacist or healthcare provider

What are the ingredients in this Medicated Cream?

Active Ingredients:

- **Lignocaine:** Local anesthetic for numbing pain
- **Metronidazole:** Antibacterial agent effective against anaerobic bacteria
- **Sucralfate:** Healing agent that forms a protective barrier and promotes tissue regeneration

Inactive Ingredients:

- Cream base

Note: This leaflet is for informational purposes only and does not replace medical advice. Always consult your healthcare provider for any concerns regarding your condition or treatment.