

Patient & Product Information

(Chlorhexidine Gluconate 1.0% w/w, Metronidazole 1.0% w/w & Lignocaine 2.0% w/w gel)
Antiseptic, Antibacterial & Anesthetic Oral Gel

Active Ingredients:

Chlorhexidine Gluconate 1.0% w/w
Metronidazole 1.0% w/w
Lignocaine (Lidocaine)..... 2.0% w/w

Read this leaflet before using the product. This information does not replace medical advice from your doctor or dentist.

What is this Oral Gel used for?

This triple-action medicated gel is indicated for the treatment and relief of:

- Gingivitis and periodontitis (gum infections)
- Oral ulcers, mouth sores, and oral cavity infections
- Pain relief in inflamed or infected areas of the mouth
- Post-dental procedures (e.g., scaling, extraction) to prevent infection and promote healing

How does it work?

- Chlorhexidine Gluconate – A broad-spectrum antiseptic effective against plaque and oral bacteria
- Metronidazole – An antibacterial and antiprotozoal agent that targets anaerobic infections
- Lignocaine – A fast-acting local anesthetic that relieves pain and irritation

Who should NOT use this Gel?

Do not use if you:

- Are allergic to chlorhexidine, metronidazole, lignocaine, or any of the gel's ingredients
- Have a history of lidocaine sensitivity or metronidazole intolerance

Area child under 12 years, unless directed by a healthcare provider?

Precautions:

- Before using, inform your doctor or dentist if you:
- Are pregnant or breastfeeding
- Have a history of seizures (due to metronidazole)
- Are using other topical or oral medications
- Have liver problems, as metronidazole is metabolized in the liver

How to Use this Oral Gel:

- Wash hands before application
- Apply a small quantity (pea-sized) of the gel directly to the affected area using a clean fingertip or cotton bud
- Use 2 to 3 times daily or as prescribed
- Do not eat or drink for 30 minutes after application to allow proper absorption
- Avoid swallowing the gel
- Do not rinse your mouth immediately after application

Possible Side Effects:

Common:

- Temporary numbness, tingling, or burning sensation
- Mild taste alteration
- Local irritation or redness

Rare but serious:

- Allergic reaction: rash, itching, swelling, or difficulty breathing

- Prolonged numbness or oral discomfort
- If you notice any of the above or symptoms worsen, stop use and consult your healthcare provider.

How should I store this Cream?

- Store at room temperature (20°C to 25°C or 68°F to 77°F)
- Do not freeze
- Keep the tube tightly closed after use
- Keep out of reach of children

General Safety Advice:

- For external oral use only – do not swallow
- Do not use for longer than prescribed duration
- Avoid use near eyes or nose – rinse thoroughly with water if contact occurs
- Do not share with others
- Use under dental or medical supervision for long-term treatment

What are the ingredients in this Cream?**Active Ingredients:**

- **Chlorhexidine Gluconate** – A broad-spectrum antiseptic effective against plaque and oral bacteria
- **Metronidazole** – An antibacterial and antiprotozoal agent that targets anaerobic infections
- **Lignocaine** – A fast-acting local anesthetic that relieves pain and irritation

Inactive Ingredients:

- Gel base

Note: This leaflet is for informational purposes only and does not replace medical advice. Always consult your healthcare provider for any concerns regarding your condition or treatment.

For more information, contact Peak Life Innovations Pvt. Ltd. at (Peaklifeinnovations@gmail.com)