

Patient & Product Information

(Clobetasol Propionate (0.05% w/w), Neomycin Sulphate 0.5% (equivalent to Neomycin base 0.35%) & Miconazole Nitrate (2% w/w) Cream) Medicated **Cream**

Active Ingredients:

Clobetasol Propionate 0.05% w/w

Neomycin Sulphate 0.5% w/w (equivalent to Neomycin base 0.35% w/w)

Miconazole Nitrate 2.0% w/w

Read this Patient Information before using this cream and each time you get a refill. This leaflet does not replace the advice of your healthcare provider.

What is this Cream used for?

This combination cream is used for the topical treatment of mixed skin infections, particularly when inflammation, bacterial, and fungal infections are present.

Functions:

- **Clobetasol Propionate:** Potent corticosteroid to reduce redness, itching, and swelling
- **Neomycin Sulphate:** Broad-spectrum antibiotic for bacterial skin infections
- **Miconazole Nitrate:** Antifungal agent for fungal infections such as ringworm, athlete's foot, and candidiasis

Who should not use this Cream?

Do not use if you:

- Are allergic to any of the ingredients
- Have viral skin infections (e.g., herpes simplex, chickenpox)
- Are pregnant or breastfeeding, unless directed by your healthcare provider
- Have extensive skin damage or open wounds

Who can use this Cream?

Adults and Children 12 Years & Older:

- Safe under medical supervision
- Apply only on affected areas in thin layers

Children under 12 Years:

- Use only if prescribed
- Avoid long-term or large-area application

What should I tell my healthcare provider before using this Cream?

Before using this product, inform your provider if you:

- Have any known drug allergies
- Are using other skin medications
- Are pregnant or breastfeeding
- Have skin sensitivity or previous reactions to topical antibiotics/steroids

How should I use this Cream?

- Clean and dry the affected area thoroughly
- Apply a thin layer to the affected skin 1–2 times daily
- Do not use occlusive dressings unless directed
- Wash hands after applying
- Avoid contact with eyes, nose, mouth, and broken skin

Possible Side Effects:

Common:

- Mild burning, stinging, or redness
- Temporary skin dryness or irritation

Rare but Serious:

- Allergic reactions (rash, swelling, difficulty breathing)
- Prolonged use may cause skin thinning, pigmentation changes
- Overuse of Neomycin can lead to skin sensitization

How should I store this Cream?

Storage Instructions:

- Store below 25°C (77°F)
- Do not freeze
- Keep tightly closed and out of reach of children

General Precautions:

- Do not use longer than 2 weeks without medical supervision
- Do not apply to large surface areas or under bandages unless advised
- Avoid prolonged use on the face or genitals
- For external use only

What are the ingredients in this Cream?

Active Ingredients:

- **Clobetasol Propionate** – Corticosteroid (anti-inflammatory)
- **Neomycin Sulfate** – Antibiotic (broad-spectrum, bacterial infections)
- **Miconazole Nitrate** – Antifungal (Candida, dermatophytes)

Inactive Ingredients:

- Cream base, Emulsifiers, Preservative

Note: This leaflet is for informational purposes only and does not replace medical advice. Always consult your healthcare provider for any concerns regarding your condition or treatment.